

Bay Area Orthopedic Surgery and Sports Medicine

100 Hospital Drive
Suite 303
Vallejo, CA 94939
Ph: 707-645-7210 fax: 707-645-7249
www.baosurgery.com

Teodoro P. Nissen, MD

*Fellowship Trained
Sports Medicine and Sports Trauma
Reconstructive Joint Surgery*

Elizabeth J. Milne, MHS, PA-C

*Physician Assistant – Certified
Orthopedics and Pain Management*

Associate Physicians:

Claudio Palma, MD

*SOMA Orthopedics
Fellowship Trained
Pain Management
415-642-7070*

Dan Birkbeck, MD

*Napa Valley Orthopedics
Fellowship Trained
Hand and Microvascular Surgery
707-647-2053*

Jason Huffman, MD

*Napa Valley Orthopedics
Fellowship Trained
Spine Surgery
707-647-2053*

Dislocated Shoulder

The shoulder joint is the body's most mobile joint. It can turn in many directions. But, this advantage also makes the shoulder an easy joint to dislocate. A partial dislocation (subluxation) means the head of the upper arm bone (humerus) is partially out of the socket (glenoid). A complete dislocation means it is all the way out of the socket. Both partial and complete dislocation cause pain and unsteadiness in the shoulder.

Symptoms

Symptoms to look for include:

- Swelling
- Numbness
- Weakness
- Bruising

Sometimes dislocation may tear ligaments or tendons in the shoulder or damage nerves.

The shoulder joint can dislocate forward, backward, or downward. A common type of shoulder dislocation is when the shoulder slips forward (anterior instability). This means the upper arm bone moved forward and down out of its joint. It may happen when the arm is put in a throwing position.

Diagnosis

The muscles may have spasms from the disruption, and this can make it hurt more. When the shoulder dislocates time and again, there is shoulder instability.

The doctor will examine the shoulder and may order an X-ray. It is important that the doctor know how the dislocation happened and whether the shoulder had ever been dislocated before.

Treatment

The doctor will place the ball of the upper arm bone (humerus) back into the joint socket. This process is called closed reduction. Severe pain stops almost immediately once the shoulder joint is back in place.

Rehabilitation

Your doctor may immobilize the shoulder in a sling or other device for several weeks following treatment. Plenty of rest is needed. The sore area can be iced 3 to 4 times a day.

After the pain and swelling go down, the doctor will prescribe rehabilitation exercises for you. These help restore the shoulder's range of motion and strengthen the muscles. Rehabilitation may also help prevent dislocating the shoulder again in the future. Rehabilitation will begin with gentle muscle toning exercises. Later, weight training can be added.

If shoulder dislocation becomes a chronic condition, a brace can sometimes help. However, if therapy and bracing fail, surgery may be needed to repair or tighten the torn or stretched ligaments that help hold the joint in place, particularly in young athletes.